

96-1-14/31

Increasing the Economic Effect of Thermal De-aeration of Feedwater
for Steam Boilers.

carrying out lime and magnesia de-silication of water at temperatures of 20 - 25 °C should be investigated. If the conditions are favourable, delivery of water from the lime-cationite installations to the turbine condensers should be tried.

There are 1 figure and 3 tables.

ASSOCIATION: TEP - VTI

AVAILABLE: Library of Congress.

Card 3/3

GUREVICH, L.S., montazhnik; ZIMENKO, N.A., montazhnik.

Light-duty screw jack with a capacity of 10 tons. Suggested by L.S. Gurevich, N.A. Zimenko. Rats. i izobr. predl. v stroi. no. 15:50 '60.
(MIRA 13:9)

1. Po materialam Zaporozhskogo stroitel'no-montazhnogo upravleniya No. 203 tresta Metallurgmontazh Ministerstva stroitel'stva USSR.
(Lifting jacks)

PROKHOROV, F.G., kand.tekhn.nauk; GUREVICH, L.S., inzh.

Features of the water treating systems of electric power plants
being constructed in the present seven-year plan.

Teploenergetika 8 no.4:3-6 Ap '61. (MIRA 14:8)

1. Ministerstvo stroitel'stva elektrostantsiy, Vsesoyuznyy
gosudarstvennyy proyektnyy institut Teploelektroproyekt.
(Electric power plants)
(Feed-water purification)

GUREVICH, Leonid Samoylovich, zhurnal'ist(Riga); NOSOV, Sergey
Petrovich, nauchnyy sotr.; GUDKOVA, N., red.; SEMENOVA, O.,
tekhn. red.

[Army Commander Aleksandr Stepin']Komandarm Aleksandr Stepin'.
Moskva, Gospolitizdat, 1962. 52 p. (MIRA 15:12)

1. Gosudarstvennyy arkhiv Sovetskoy Armii (for Nosov).
(Stepin', Aleksandr Karlovich, d.1920)

SHEYBER, B.P., kand. tekhn. nauk; GUREVICH, L.S., inzh.

Set of the EO-1 equipment for preliminary and subsequent
bituminization. Transp. stroi. 15 no.2:52-53 F '65. (MIRA 18:3)

PROKHOROV, F.G., kand.tekhn.nauk; GUREVICH, L.S., inzh.

Comparison of water conditions and means for water treatment in
block-type state regional electric power plants with different
boiler systems. Teploenergetika 12 no.10:2-8 0 '65.

(MIRA 18:10)

1. Vsesoyuznyy nauchno-issledovatel'skiy teplotekhnicheskiy institut
i Vsesoyuznyy gosudarstvennyy proyektnyy institut "Teploelektroproyekt".

GUREVICH, L. SH.

USSR/Chemistry - Synthesis

Card 1/1 Pub. 151 - 34/38

Authors : Khaletskiy, A. M., and Gurevich, L. Sh.

Title : Synthesis of ethyl ether and amide of α_1 -amino nicotinic acid.

Periodical : Zhur. ob. khim. 24/2, 369-372, Feb 1954

Abstract : The synthesis of ethyl ether and amide of α_1 -amino nicotinic acid from technical anabasine-sulfate was investigated. The ether was obtained through oxidation of α_1 -chloroanabasine followed by amination and etherification of the formed α_1 -amino nicotinic acid. Treatment of the ether with ammonia under pressure leads to the formation of amide of α_1 -amino nicotinic acid. The anesthetic effect of the ether was established through pharmacological research. Three USSR references (1931-1941).

Institution: The Chemical-Pharmaceutical Institute, Leningrad

Submitted : September 26, 1953

GUREVICH, L. Sh.

USSR.

✓ Synthesis of the ethyl ester and amide of α -aminocaproic acid. A. M. Khaletskii and L. Sh. Gurevich. *J. Gen. Chem. U.S.S.R.* 24, 377-0 (1954) (Eng. transl.).
Sep. C.A. 49, 4649h. H. L. H.

GUREVICH, L., kand. tekhn. nauk; TURCHIKHIN, E., kand. tekhn. nauk

Using colored materials in constructing pavements. Zhil.-kon. khoz.
9 no.9:16-17 '59. (MIRA 13:2)
(Pavements)

RODINOV, V.P., assistant; GUREVICH, L.V., tech.

Balancing of engines in assembly at the Automobile Repair Plant
No.4. Izv. vys. ucheb. zav.; mashinostr. no.3:100-102 '64.
(MIRA 17:7)

1. Morskoye vyssheye tekhnicheskoye uchilishche
imeni Baumana.

L 39673-65

ACCESSION NR: AP5010475

UR/0294/65/003/002/0318/0321

AUTHOR: Ryabova, V. G.; Gurvich, L. V.

TITLE: Study of metal-hydroxyl bond energies in CaOH, SrOH, and BaOH molecules

SOURCE: Teplofizika vysokikh temperatur, v. 3, no. 2, 1969, 318-321

TOPIC TAGS: combustion, hydrogen oxygen, metallized fuel, metal combustion, flame spectroscopy

ABSTRACT: The energies of the hydroxyl-metal bonds in CaOH, SrOH, and BaOH, which may be present together with the oxides in hydrogen-air flames, were determined spectroscopically by studying the reaction of Ca, Sr, and Ba with combustion products of hydrogen-air flames having different compositions. The metals were introduced as SrCl_2 , BaCl_2 and $\text{Ca}(\text{COO})_2$ solutions into the flame of a Meker burner. The partial pressures of the metals were determined from the spectral line intensities measured at flame temperatures of 1760—2160K. The experiments yielded bond energies of about 101, 97, and 112 kcal/mol for Ca, Sr, and Ba, respectively. These values indicate, as was shown by thermodynamic calculations, that a considerable amount of the metals must be present in rich hydrogen-air flames as Me-OH, formed by the reaction $\text{Me} + \text{H}_2\text{O} = \text{MeOH} + \text{H}$. The bond energies of MeO were

Card 1/2

L 39673-65

ACCESSION NR: AP5010475

recalculated from previous experimental data obtained with acetylene-air, acetylene-oxygen, hydrogen-oxygen, and monoxide-oxygen flames. Orig. art. has: 2 tables.

[FV]

ASSOCIATION: Nauchno-issledovatel'skiy institut vysokikh temperatur (Scientific Research Institute of High Temperatures)

SUBMITTED: 30Oct64

ENCL: 00

SUB CODE: FT

NO REF SOV: 012

OTHER: 004

ATD PRESS: 3229

Card 2/2

GUREVICH, L. V. Cand. Tech. Sci.

Dissertation: "Premises For Solution of the Problem of Designing Paired Highways and Railroads." Moscow Inst of Engineers of Municipal Building, 11 Feb 47.

SO: Vechernyaya Moskva, Feb, 1947 (Project #17836)

SECRET
GUREVICH, L.V.; YAMPOL'SKAYA, T.G.; MURZAYEVA, L.B.; KHRUNOV, N.P., redaktor;
OTOCHIEVA, M.A., redaktor; PETROVSKAYA, Ye., tekhnicheskii redaktor

[Road traffic signs] Dorozhnye signal'nye znaki. Moskva, Izd-vo
Ministerstva kommunal'nogo khoziaistva RSFSR, 1955. 46 p.
(Traffic regulations) (MLRA 9:2)

Gurevich, L.V.
USSR/Optics - Physiological Optics

K-9

Abs Jour : Referat Zhur - Fizika, No 5, 1957, 13180

Author : Gurtovoy, G.K., Gurevich, L.V., Murzayeva, L.B.,
Seletskaya, L.I., Yampol'skaya, T.G.

Inst : -

Title : Investigation of the Laws of Color and Three Dimensional
Visions and Their Use for Increasing the Effectiveness of
Road Signals.

Orig Pub : Tr. In-ta biol. fiz. AN SSSR, 1955, 1, 136-157

Abstract : Starting with the premise that the problems of visibility
of road signals are insufficiently well developed, the
authors have undertaken an extensive investigation of the
influence of such factors, as the shape of the signs, the
combination of colors of the image on the sign and of the
background, the dimension of the sign, and its illumina-
tion. As a total the following recommendations were made:
(1) With respect to the shape -- rectangle (1:4 to 1:10),

Card 1/2

00513R000617420006-5

SOSYANTS, V.G.; OVECHNIKOV, Ye.V.; GUREVICH, L.V.; LESEVITSKIY, N.N.;
BASHKIROV, L.G., redaktor; KONYASHINA, A., tekhnicheskii redaktor

[Construction of trolley tracks with concrete foundations] Kon-
struktsii tramvaynykh putei s betonnyimi osnovaniyami. Moskva,
Izd-vo Ministerstva kommunal'nogo khoziaistva RSFSR, 1956. 52 p.
(Street railways) (MLRA 9:11)

GUREVICH, I. I., KURATOV, I. A., SHTEKLYA, I. I., KUMOLEV, T. Ya., GUR'VA, G. F.

Investigation of the Interrelationships Underlying Color and Space Vision and
Application of Results Obtained in Increasing the Effectiveness of Road Signs

Trudy Instituta Biologicheskoy Fiziki, No 1, 1956
S916, 5 Mar 1956, p 49

GUREVICH, L.V.

Multiple-seated slabs in road construction. Gor. khoz. Mosk.
30 no.9:39 S '56. (MLRA 9:12)

1. Starshiy nauchnyy sotrudnik Akademii kommunal'nogo
khozaystva.
(Concrete slabs) (Road construction)

GUREVICH, L.V.; DITERIKHS, N.D.; LUCHAY, G.A.; NIKOL'SKAYA, N.Ye.

Using plastics in the rolling stock and in the electric power
supply of public transportation. Sbor.nauch.rab.AKKH no.13:192-
202 '62. (MIRA 16:4)

(Plastics)

(Local transit—Equipment and supplies)

SOSYANTS, V.G., inzh.; YUDIN, V.A., kand. tekhn.nauk; KNORRE, V.E., inzh.; LANTSEBERG, Yu.S., inzh.; DAVIDYANTS, N.M., inzh.; GEZENTSVEY, L.B., kand. tekhn. nauk; YEGOROV, P.A., inzh.; FAYNBERG, E.S., inzh.; BAGDASAROV, S.M., inzh.; GUREVICH, L.V., kand. tekhn. nauk; CHERNYSHOV, B.G., inzh.; GADZHINSKIY, T.G., inzh.; ZASOV, I.A., kand. tekhn.nauk; BALOVNEV, V.I., kand. tekhn.nauk; GIBSHMAN, Ye.Ye., prof., red.; DZHUNKOVSKIY, N.N., prof., red.; BOLOTINA, A.V., red. izd-va; LELYUKHIN, A.A., tekhn. red.

[Manual for the design, construction, and maintenance of urban roads, bridges, and hydrotechnical structures]
Spravochnik po proektirovaniu, stroitel'stvu i ekspluatatsii gorodskikh dorog, mostov i gidrotekhnicheskikh sooruzhenii. Red. kol.E.E.Gibshman, N.N.Dzhunkovskii, P.A. Egorov. Moskva, Izd-vo M-va kommun.khoz.RSFSR. Vol.3. [Roads] Dorogi. 1963. 814 p. (MIRA 16:7)
(Roads)

GIREVICH, L. V., kand.tekhn.nauk; SOSKIN, G. M., kand.tekhn.nauk

Precast street and sidewalk pavement. Nov.tekh.zhil.-kom.khoz.:
Gor.dor.-most.khoz. i transp. no. 2:5-9 '63. (MIRA 17:5)

GUREVICH, L. V., kand.tekhn.nauk; GURFINK, T. Sh., inzh.; SOSEKIN, G. M.,
kand. tekhn. nauk

Landscaping and problems in designing city streets. Nov.tekh.
zhil.-kom.khoz.:Gor.dor.-most.khoz. i transp. no. 2:9-12 '63.
(MIRA 17:5)

1ST AND 2ND ORDERS										PROCESSING AND PROPERTY INDEX									
<p><i>BC</i></p> <p>Polymerization processes induced by mitogenetic radiation. A. GUNVIRSON and L. GUNVIRSON (Acta Physicochim. U.R.S.S., 1939, 10, 711-718).—When a freshly prepared 10% peptone (I) solution is irradiated with mitogenetic radiation a substance is produced which reacts with gastric juice, whilst the original (I) does not. If the peptone solution is shaken with kaolin, the latter separated, and the solution is then irradiated, the same effect is obtained, but if the shaking with kaolin is carried out after the irradiation, there is no effect. The absorption of irradiated (I) in the range 2800—2400 Å. is considerably > that of the original substance. These phenomena point to the formation of a polymeride under the influence of the radiation. The concn. of the polymeride, within certain limits, or that of the original peptone solution. The polymerization of a dipeptide (glycylglycine) by mitogenetic radiation was also investigated. After irradiation the product had</p> <p>no effect on gastric juice, but a positive effect was obtained after the addition of a small amount of serum-albumin.</p> <p style="text-align: right;">A. J. M.</p>										<p>171</p>									
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>										<p>ADMIN. NUMBER</p>									
<p>1ST AND 2ND ORDERS</p>										<p>1ST AND 2ND ORDERS</p>									

BC

Explanation of mitogenetic radiation as "sensitized fluorescence." A. GURVITSKY and L. GUMVITSKY (Acta Physicochim. U.R.S.S., 1939, 10, 719-724).—Frankenburger's theory, that energy made available in org. and particularly fermentation, reactions taking place by intermediate formation of atoms and radicals, can be used to excite mol. of other substances present giving rise to mitogenetic radiation, has been confirmed experimentally. When glycine, previously irradiated, was mixed with glucose undergoing fermentation, the characteristic glucose bands at 1900—1905 and 1915—1920 Å. were excited. No fermentation product could act as the receptor in this case. Na⁺ and Cl⁻ can act as receptors. The bearing of these results on mitogenetic spectral analysis is discussed. A. J. M.

ASAC SLA METALLURGICAL LITERATURE CLASSIFICATION

1900 1910 1920 1930 1940 1950 1960 1970 1980 1990 2000 2010 2020 2030 2040 2050 2060 2070 2080 2090 2100 2110 2120 2130 2140 2150 2160 2170 2180 2190 2200 2210 2220 2230 2240 2250 2260 2270 2280 2290 2300 2310 2320 2330 2340 2350 2360 2370 2380 2390 2400 2410 2420 2430 2440 2450 2460 2470 2480 2490 2500 2510 2520 2530 2540 2550 2560 2570 2580 2590 2600 2610 2620 2630 2640 2650 2660 2670 2680 2690 2700 2710 2720 2730 2740 2750 2760 2770 2780 2790 2800 2810 2820 2830 2840 2850 2860 2870 2880 2890 2900 2910 2920 2930 2940 2950 2960 2970 2980 2990 3000 3010 3020 3030 3040 3050 3060 3070 3080 3090 3100 3110 3120 3130 3140 3150 3160 3170 3180 3190 3200 3210 3220 3230 3240 3250 3260 3270 3280 3290 3300 3310 3320 3330 3340 3350 3360 3370 3380 3390 3400 3410 3420 3430 3440 3450 3460 3470 3480 3490 3500 3510 3520 3530 3540 3550 3560 3570 3580 3590 3600 3610 3620 3630 3640 3650 3660 3670 3680 3690 3700 3710 3720 3730 3740 3750 3760 3770 3780 3790 3800 3810 3820 3830 3840 3850 3860 3870 3880 3890 3900 3910 3920 3930 3940 3950 3960 3970 3980 3990 4000 4010 4020 4030 4040 4050 4060 4070 4080 4090 4100 4110 4120 4130 4140 4150 4160 4170 4180 4190 4200 4210 4220 4230 4240 4250 4260 4270 4280 4290 4300 4310 4320 4330 4340 4350 4360 4370 4380 4390 4400 4410 4420 4430 4440 4450 4460 4470 4480 4490 4500 4510 4520 4530 4540 4550 4560 4570 4580 4590 4600 4610 4620 4630 4640 4650 4660 4670 4680 4690 4700 4710 4720 4730 4740 4750 4760 4770 4780 4790 4800 4810 4820 4830 4840 4850 4860 4870 4880 4890 4900 4910 4920 4930 4940 4950 4960 4970 4980 4990 5000 5010 5020 5030 5040 5050 5060 5070 5080 5090 5100 5110 5120 5130 5140 5150 5160 5170 5180 5190 5200 5210 5220 5230 5240 5250 5260 5270 5280 5290 5300 5310 5320 5330 5340 5350 5360 5370 5380 5390 5400 5410 5420 5430 5440 5450 5460 5470 5480 5490 5500 5510 5520 5530 5540 5550 5560 5570 5580 5590 5600 5610 5620 5630 5640 5650 5660 5670 5680 5690 5700 5710 5720 5730 5740 5750 5760 5770 5780 5790 5800 5810 5820 5830 5840 5850 5860 5870 5880 5890 5900 5910 5920 5930 5940 5950 5960 5970 5980 5990 6000 6010 6020 6030 6040 6050 6060 6070 6080 6090 6100 6110 6120 6130 6140 6150 6160 6170 6180 6190 6200 6210 6220 6230 6240 6250 6260 6270 6280 6290 6300 6310 6320 6330 6340 6350 6360 6370 6380 6390 6400 6410 6420 6430 6440 6450 6460 6470 6480 6490 6500 6510 6520 6530 6540 6550 6560 6570 6580 6590 6600 6610 6620 6630 6640 6650 6660 6670 6680 6690 6700 6710 6720 6730 6740 6750 6760 6770 6780 6790 6800 6810 6820 6830 6840 6850 6860 6870 6880 6890 6900 6910 6920 6930 6940 6950 6960 6970 6980 6990 7000 7010 7020 7030 7040 7050 7060 7070 7080 7090 7100 7110 7120 7130 7140 7150 7160 7170 7180 7190 7200 7210 7220 7230 7240 7250 7260 7270 7280 7290 7300 7310 7320 7330 7340 7350 7360 7370 7380 7390 7400 7410 7420 7430 7440 7450 7460 7470 7480 7490 7500 7510 7520 7530 7540 7550 7560 7570 7580 7590 7600 7610 7620 7630 7640 7650 7660 7670 7680 7690 7700 7710 7720 7730 7740 7750 7760 7770 7780 7790 7800 7810 7820 7830 7840 7850 7860 7870 7880 7890 7900 7910 7920 7930 7940 7950 7960 7970 7980 7990 8000 8010 8020 8030 8040 8050 8060 8070 8080 8090 8100 8110 8120 8130 8140 8150 8160 8170 8180 8190 8200 8210 8220 8230 8240 8250 8260 8270 8280 8290 8300 8310 8320 8330 8340 8350 8360 8370 8380 8390 8400 8410 8420 8430 8440 8450 8460 8470 8480 8490 8500 8510 8520 8530 8540 8550 8560 8570 8580 8590 8600 8610 8620 8630 8640 8650 8660 8670 8680 8690 8700 8710 8720 8730 8740 8750 8760 8770 8780 8790 8800 8810 8820 8830 8840 8850 8860 8870 8880 8890 8900 8910 8920 8930 8940 8950 8960 8970 8980 8990 9000 9010 9020 9030 9040 9050 9060 9070 9080 9090 9100 9110 9120 9130 9140 9150 9160 9170 9180 9190 9200 9210 9220 9230 9240 9250 9260 9270 9280 9290 9300 9310 9320 9330 9340 9350 9360 9370 9380 9390 9400 9410 9420 9430 9440 9450 9460 9470 9480 9490 9500 9510 9520 9530 9540 9550 9560 9570 9580 9590 9600 9610 9620 9630 9640 9650 9660 9670 9680 9690 9700 9710 9720 9730 9740 9750 9760 9770 9780 9790 9800 9810 9820 9830 9840 9850 9860 9870 9880 9890 9900 9910 9920 9930 9940 9950 9960 9970 9980 9990 10000 10010 10020 10030 10040 10050 10060 10070 10080 10090 10100 10110 10120 10130 10140 10150 10160 10170 10180 10190 10200 10210 10220 10230 10240 10250 10260 10270 10280 10290 10300 10310 10320 10330 10340 10350 10360 10370 10380 10390 10400 10410 10420 10430 10440 10450 10460 10470 10480 10490 10500 10510 10520 10530 10540 10550 10560 10570 10580 10590 10600 10610 10620 10630 10640 10650 10660 10670 10680 10690 10700 10710 10720 10730 10740 10750 10760 10770 10780 10790 10800 10810 10820 10830 10840 10850 10860 10870 10880 10890 10900 10910 10920 10930 10940 10950 10960 10970 10980 10990 11000 11010 11020 11030 11040 11050 11060 11070 11080 11090 11100 11110 11120 11130 11140 11150 11160 11170 11180 11190 11200 11210 11220 11230 11240 11250 11260 11270 11280 11290 11300 11310 11320 11330 11340 11350 11360 11370 11380 11390 11400 11410 11420 11430 11440 11450 11460 11470 11480 11490 11500 11510 11520 11530 11540 11550 11560 11570 11580 11590 11600 11610 11620 11630 11640 11650 11660 11670 11680 11690 11700 11710 11720 11730 11740 11750 11760 11770 11780 11790 11800 11810 11820 11830 11840 11850 11860 11870 11880 11890 11900 11910 11920 11930 11940 11950 11960 11970 11980 11990 12000 12010 12020 12030 12040 12050 12060 12070 12080 12090 12100 12110 12120 12130 12140 12150 12160 12170 12180 12190 12200 12210 12220 12230 12240 12250 12260 12270 12280 12290 12300 12310 12320 12330 12340 12350 12360 12370 12380 12390 12400 12410 12420 12430 12440 12450 12460 12470 12480 12490 12500 12510 12520 12530 12540 12550 12560 12570 12580 12590 12600 12610 12620 12630 12640 12650 12660 12670 12680 12690 12700 12710 12720 12730 12740 12750 12760 12770 12780 12790 12800 12810 12820 12830 12840 12850 12860 12870 12880 12890 12900 12910 12920 12930 12940 12950 12960 12970 12980 12990 13000 13010 13020 13030 13040 13050 13060 13070 13080 13090 13100 13110 13120 13130 13140 13150 13160 13170 13180 13190 13200 13210 13220 13230 13240 13250 13260 13270 13280 13290 13300 13310 13320 13330 13340 13350 13360 13370 13380 13390 13400 13410 13420 13430 13440 13450 13460 13470 13480 13490 13500 13510 13520 13530 13540 13550 13560 13570 13580 13590 13600 13610 13620 13630 13640 13650 13660 13670 13680 13690 13700 13710 13720 13730 13740 13750 13760 13770 13780 13790 13800 13810 13820 13830 13840 13850 13860 13870 13880 13890 13900 13910 13920 13930 13940 13950 13960 13970 13980 13990 14000 14010 14020 14030 14040 14050 14060 14070 14080 14090 14100 14110 14120 14130 14140 14150 14160 14170 14180 14190 14200 14210 14220 14230 14240 14250 14260 14270 14280 14290 14300 14310 14320 14330 14340 14350 14360 14370 14380 14390 14400 14410 14420 14430 14440 14450 14460 14470 14480 14490 14500 14510 14520 14530 14540 14550 14560 14570 14580 14590 14600 14610 14620 14630 14640 14650 14660 14670 14680 14690 14700 14710 14720 14730 14740 14750 14760 14770 14780 14790 14800 14810 14820 14830 14840 14850 14860 14870 14880 14890 14900 14910 14920 14930 14940 14950 14960 14970 14980 14990 15000 15010 15020 15030 15040 15050 15060 15070 15080 15090 15100 15110 15120 15130 15140 15150 15160 15170 15180 15190 15200 15210 15220 15230 15240 15250 15260 15270 15280 15290 15300 15310 15320 15330 15340 15350 15360 15370 15380 15390 15400 15410 15420 15430 15440 15450 15460 15470 15480 15490 15500 15510 15520 15530 15540 15550 15560 15570 15580 15590 15600 15610 15620 15630 15640 15650 15660 15670 15680 15690 15700 15710 15720 15730 15740 15750 15760 15770 15780 15790 15800 15810 15820 15830 15840 15850 15860 15870 15880 15890 15900 15910 15920 15930 15940 15950 15960 15970 15980 15990 16000 16010 16020 16030 16040 16050 16060 16070 16080 16090 16100 16110 16120 16130 16140 16150 16160 16170 16180 16190 16200 16210 16220 16230 16240 16250 16260 16270 16280 16290 16300 16310 16320 16330 16340 16350 16360 16370 16380 16390 16400 16410 16420 16430 16440 16450 16460 16470 16480 16490 16500 16510 16520 16530 16540 16550 16560 16570 16580 16590 16600 16610 16620 16630 16640 16650 16660 16670 16680 16690 16700 16710 16720 16730 16740 16750 16760 16770 16780 16790 16800 16810 16820 16830 16840 16850 16860 16870 16880 16890 16900 16910 16920 16930 16940 16950 16960 16970 16980 16990 17000 17010 17020 17030 17040 17050 17060 17070 17080 17090 17100 17110 17120 17130 17140 17150 17160 17170 17180 17190 17200 17210 17220 17230 17240 17250 17260 17270 17280 17290 17300 17310 17320 17330 17340 17350 17360 17370 17380 17390 17400 17410 17420 17430 17440 17450 17460 17470 17480 17490 17500 17510 17520 17530 17540 17550 17560 17570 17580 17590 17600 17610 17620 17630 17640 17650 17660 17670 17680 17690 17700 17710 17720 17730 17740 17750 17760 17770 17780 17790 17800 17810 17820 17830 17840 17850 17860 17870 17880 17890 17900 17910 17920 17930 17940 17950 17960 17970 17980 17990 18000 18010 18020 18030 18040 18050 18060 18070 18080 18090 18100 18110 18120 18130 18140 18150 18160 18170 18180 18190 18200 18210 18220 18230 18240 18250 18260 18270 18280 18290 18300 18310 18320 18330 18340 18350 18360 18370 18380 18390 18400 18410 18420 18430 18440 18450 18460 18470 18480 18490 18500 18510 18520 18530 18540 18550 18560 18570 18580 18590 18600 18610 18620 18630 18640 18650 18660 18670 18680 18690 18700 18710 18720 18730 18740 18750 18760 18770 18780 18790 18800 18810 18820 18830 18840 18850 18860 18870 18880 18890 18900 18910 18920 18930 18940 18950 18960 18970 18980 18990 19000 19010 19020 19030 19040 19050 19060 19070 19080 19090 19100 19110 19120 19130 19140 19150 19160 19170 19180 19190 19200 19210 19220 19230 19240 19250 19260 19270 19280 19290 19300 19310 19320 19330 19340 19350 19360 19370 19380 19390 19400 19410 19420 19430 19440 19450 19460 19470 19480 19490 19500 19510 19520 19530 19540 19550 19560 19570 19580 19590 19600 19610 19620 19630 19640 19650 19660 19670 19680 19690 19700 19710 19720 19730 19740 19750 19760 19770 19780 19790 19800 19810 19820 19830 19840 19850 19860 19870 19880 19890 19900 19910 19920 19930 19940 19950 19960 19970 19980 19990 20000 20010 20020 20030 20040 20050 20060 20070 20080 20090 20100 20110 20120 20130 20140 20150 20160 20170 20180 20190 20200 20210 20220 20230 20240 20250 20260 20270 20280 20290 20300 20310 20320 20330 20340 20350 20360 20370 20380 20390 20400 20410 20420 20430 20440 20450 20460 20470 20480 20490 20500 20510 20520 20530 20540 20550 20560 20570 20580 20590 20600 20610 20620 20630 20640 20650 20660 20670 20680 20690 20700 20710 20720 20730 20740 20750 20760 20770 20780 20790 20800 20810 20820 20830 20840 20850 20860 20870 20880 20890 20900 20910 20920 20930 20940 20950 20960 20970 20980 20990 21000 21010 21020 21030 21040 21050 21060 21070 21080 21090 21100 21110 21120 21130 21140 21150 21160 21170 21180 21190 21200 21210 21220 21230 21240 21250 21260 21270 21280 21290 21300 21310 21320 21330 21340 21350 21360 21370 21380 21390 21400 21410 21420 21430 21440 21450 21460 21470 21480 21490 21500 21510 21520 21530 21540 21550 21560 21570 21580 21590 21600 21610 21620 21630 21640 21650 21660 21670 21680 21690 21700 21710 21720 21730 21740 21750 21760 21770 21780 21790 21800 21810 21820 21830 21840 21850 21860 21870 21880 21890 21900 21910 21920 21930 21940 21950 21960 21970 21980 21990 22000 22010 22020 22030 22040 22050 22060 22070 22080 22090 22100 22110 22120 22130 22140 22150 22160 22170 22180 22190 22200 22210 22220 22230 22240 22250 22260 22270 22280 22290 22300 22310 22320 22330 22340 22350 22360 22370 22380 22390 22400 22410 22420 22430 22440 22450 22460 22470 22480 22490 22500 22510 22520 22530 22540 22550 22560 22570 22580 22590 22600 22610 22620 22630 22640 22650 22660 22670 22680 22690 22700 22710 22720 22730 22740 22750 22760 22770 22780 22790 22800 22810 22820 22830 22840 22850 22860 22870 22880 22890 22900 22910 22920 22930 22940 22950 22960 22970 22980 22990 23000 23010 23020 23030 23040 23050 23060 23070 23080 23090 23100 23110 23120 23130 23140 23150 23160 23170 23180 23190 23200 23210 23220 23230 2324

3C

Processes and Properties Index

Polymerization processes induced by ultraviolet radiation. II. Influence on auto-oxidation and production of a "deaminase" by irradiation. A. Gervitch and L. Gervitch (*Acta Physicochim. U.R.S.S.*, 1946, 22, 606-609; cf. A., 1936, I, 636).—The term "deaminase" is used to indicate the mode of action rather than the chemical nature of a substance (I) of high mol. wt. that is formed during the irradiation of eq. glycine. The probable sequence of reactions leading to the formation of (I) is discussed. This formation follows, but is not dependent on, the primary dissociation of the glycine mol., which is confined to the action of OH \cdot . The presence of O $_2$, though essential to the production of NH $_3$ by the action of (I) on glycine, is not needed for the formation of (I). The energy of polymerization of the active monomers that combine with mol. of (I) amounts to 71-76 kg.-cal. and may be available for the activation of the deamination process. F. L. U.

ASAC-51A METALLURGICAL LITERATURE CLASSIFICATION

SECOND HALF ONLY ONE

SCALESTONE

BC

PROCESSES AND PROPERTIES INDEX

Unstabling and inhibition of mitogenetic radiation. A. Gurvitch and I. Gurvitch (*Acta Physicochim. U.R.S.S.*, 1940, 18, 483-489).—All mol. substances which absorb in the ultra-violet can act as quenchers of every kind of mitogenetic radiation. They include not only As, I, quinine, etc., but also substances of undetermined nature that are produced by the external radiation; these latter are responsible for the rapid "fatigue" exhibited by yeast cultures, as shown by the fact that a fresh yeast culture to which a small proportion of a "fatigued" culture has been added is thereby rendered useless. Inhibitors differ from quenchers in affecting only the radiation induced by exothermic chemical reactions, including decomp. by enzymes. They are used up in the process of inhibition, and in no way disturb the course of the accompanying fermentative change. The no. of mols. of inhibitor taking an active part in inhibition is very small compared with that of the enzyme or substrate mols. taking part in the enzyme reaction. F. L. U.

ASB-11A METALLURGICAL LITERATURE CLASSIFICATION

100000 01 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

BC

Biogenetic spectrum analysis by the selective scattering method.
A. Gurvitch and L. Garryach (*Acta Physicochim. U.R.S.S.*, 1945, 20, 635-644).—In addition to selective absorption and fluorescence, the spectral composition of scattered light is determined by "chemi-fluorescence," i.e., a fluorescence related to photochemical processes occurring in the substance irradiated. The existence of chemi-fluorescence has been demonstrated by biogenetic analysis of the scattering spectrum of cholesterol, a culture of yeast on agar being used as a biological detector. The val. of the method, due to its high sensitivity, in detecting the presence of org. groups in complex mole. is illustrated (cf. A., 1946, 111, 567). C. R. H.

CA

2

Kinetics of hydrocarbon conversions in a broad range of space velocities. A. P. Balkov, L. V. Gorykh, V. V. Koro-
lov, and A. V. Frost (Moscow State Univ.). *Vestnik
Moskov. Univ.* 6, No. 2, Ser. Fiz.-Mat. i Estestv. Nauk
No. 1, 67 (1951). The applicability of the (integrated)
kinetic equation $\ln[1/(1-y)] = \beta \tau + \alpha$, characteristic
of reactions inhibited by their products, was tested in a
broad range of space velocities τ (l./l. catalyst/hr.) varied
by variation of the length of the catalyst column; y is the
degree of conversion, and α and β are consts. The test is
linearity of the plot of $\ln[1/(1-y)]$ as a function of τ .
Cracking of decahydronaphthalene (I) on activated Askan
clay (30 min runs) was found to obey this rate equation
at 300° ($\tau = 0.31-0.91$, $y = 0.51-0.20$), 400° (0.40-2.00),
at 500° ($\tau = 0.31-0.91$, $y = 0.51-0.20$), with $\alpha =$
0.51-0.181, and 500° (0.305-2.08, 0.78-0.24), with $\alpha =$
0.088, 0.129, and 0.20, resp., and $\beta = 0.35$, const. at all 3
temps. The apparent activation energy is 5.0 kcal./mole.
On an aluminosilicate cracking catalyst (SiO₂ 78.18, Al₂O₃
10.75, CaO 0.47, MgO 1.80%), at 400°, the equation holds
between $\tau = 0.6$ and 1.0, with $\alpha = 0.5$, $\beta = 0.97$. How-

ever, it fails at $\tau = 1$, the plot of $\ln[1/(1-y)]$ as a
function of τ changes from an upward sloping straight
line into a descending vertical line parallel to the axis of
ordinates, and the plot of y as a function of $1/\tau$ becomes a
straight line passing through the origin. In that range
of high τ , the reaction evidently becomes zero-order and is
described by $\tau y = 0.16$. The suspicion that this change of
kinetics might be due to a change of the Reynolds no
(which, in the cracking of I at 400°, varied from 2.8 to
26.0) was tested by the reaction of redistribution of H in an
unsatd. cracking-gasoline fraction b. 100-150° in soln. in
R₂O:Me₂CO = 1:1, followed by the decrease of the iodine
no., at 400° on the aluminosilicate catalyst in the form of
4 × 4-mm. granules in a reactor of 15-mm. diam. at a const.
Re = 17. Despite this constancy, the reaction still becomes
zero-order at high τ , above $\tau = 2.0-2.8$ ($y = 0.67-0.80$),
and then follows the equation $\tau y = 1.81$. An increase of
Re from 13 to 70 at const. τ had no effect on y . Conse-
quently, the cause of the changeover to zero order must be
chem. A clue was provided by the observation that, with 3,
the compn. of the products underwent a change at high τ ;
the compn. of the products disappeared, and the catalyzate began
to boil at 150-160°. On the basis of information in the
literature, it is assumed that the first reaction of I consists in
an isomerization to products contg. cyclopentane rings, and
that the subsequent cracking bears on such isomerization
products; this, among others, is borne out by the predom-
inance, in the gaseous cracking products, of isobutane over

over

Gurvieh, L.V.

62
 "Electronic investigation of molecular structure." I. A. V. Frost, P. A. Akishin, L. V. Gurvieh, G. A. Kurkchi, and A. A. Konstantinov (Univ. Moscow). *Vestnik Moskov. Univ.* 8, No. 12; Ser. Fiz.-Mat. i Estestv. Nauk No. 8, 85-93 (1953).—An electron-diffraction instrument for the study of any vaporizable substance is described in which a beam of electrons of 1 mm. diam., projected from an electron gun with water-cooled anode, is focused by an electromagnetic lens and diffracted by a stream of vapor of the given substance (I) onto a photographic film. The film chamber is water-cooled for high-temp. work. The vapor stream issues from a jet assembly made of Mo glass for low temps. or metal for high temps., which consists of a nozzle connected through a tube jacketed to prevent condensation to an ampul contg. the I. The position of the nozzle is adjusted with a microscope; a well is provided for the latter. A 2nd well opposite the nozzle contains liquid Na and acts as a trap for the I vapor. A sliding holder contg. a standard cryst. substance can be placed in the electron beam for calibration. This electronograph was used to det. the mol. structure of CCl_4 (II) and CdBr_2 (III) at temps. of 15 and 600°, resp. The intensities and radii of max. and min. in the diffraction patterns are tabulated and graphed. Av. values for the C—Cl and Cl—Cl distances in II are 1.756 ± 0.010 and 2.868 ± 0.015 Å., resp.; for the Cd—Br and Br—Br distances in III they are 2.35 ± 0.03 and 4.70 ± 0.03 Å., resp. The values for II and III agree within exp't. error with those of Allen and Sutton (C.A. 44, 47-46c) and Lister and S. (C.A. 36, 519), resp. J. W. L., Jr.

4

AKISHIN, P.A.; GURVICH, L.V.

Radial distribution methods in electronography and table of
values for $\sin \frac{\pi}{10} \gamma m$. Uch. zap. Mosk. un. no. 164:153-200 '53.
(MIRA 8:7)

(Electronography)

VEYTS, I.V.; GURVICH, L.V.; KOROBV, V.V.

Determination of the dissociation energy of metal oxides (SrO, CaO and MgO) by measuring the intensity of resonance lines of the metal atoms in flame spectra. Izv. AN SSSR. Ser. fiz. 19 no.1:21-22 Ja-F '55. (MIRA 8:9)

1. Institut goryuchikh iskopayemykh Akademii nauk SSSR
(Spectrum analysis) (Spectrometer)

GURVICH, L. V.

USSR/Atomic and Molecular Physics - Physics of the Molecule, D-2

Abst Journal: Referat Zhur - Fizika, No 12, 1956, 34313

Author: Veyts, I. V., Gurvich, L. V.

Institution: None

Title: Dissociation Energies of Oxides of Magnesium, Calcium, Strontium, and Barium

Original Periodical: Optika i spektroskopiya, 1956, 1, No 1, 22-33

Abstract: Based on a study of the intensities of resonant lines of metal atoms in a flame, a determination was made of the constant of equilibrium of dissociation of oxides of alkali-earth metals in flames of $C_2H_2 + \text{air}$, $C_2H_2 + O_2$, $H_2 + \text{air}$, and $H_2 + O_2$. With the aid of the so obtained values of the equilibrium constant, the energies of the dissociation of the oxides were obtained.

/ of /

- 1 -

GURVICH, L.V.: KOROBV, V.V.

Calculation of the thermodynamic functions of diatomic gases taking into account the excitation of molecular electronic states. Zhur. fiz. khim. 30 no.12:2794-2800 D'56. (MLRA 10:4)

1. Institut goryuchikh iskopayemykh, Moskva.
(Thermodynamics)

Gurvich, I.V.

USSR/ Physical Chemistry - Molecule. Chemical Bond

B-4

Abs Jour : Referat Zhur - Khimiya, No 3, 1957, 7166

Author : Veyts, I.V. and Gurvich, I.V.

Inst : Academy of Sciences USSR

Title : Dissociation Energy of AlO

Orig Pub : Dokl. AN SSSR, 1956, Vol 108, No 4, 659-661

Abstract : The energy of dissociation of AlO was calculated from the equilibrium constant for the dissociation of AlO in an oxyacetylene flame by a previously described method (RZhKhim, 1956, 74133). The partial pressure of atomic Al was calculated from the intensity of the Al(I) line at 3961.5A. A value of 133.5 ± 3 kcal/mole (5.8ev) was found for D(Al) [sic].

Card 1/1

- 13 -

GURVICH, L.V.

PRIKHOT'KO, N F

24(7) p 3 PHASE I BOOK EXPLOITATION 801/1365

L'vov. Universitet

Materialy X Vsesoyuznogo soveshchaniya po spektroskopii. t. 1: Molekulyarnaya spektroskopiya (Papers of the 10th All-Union Conference on Spectroscopy. Vol. 1: Molecular Spectroscopy) [L'vov] Izd-vo L'vovskogo univ-ta, 1957. 499 p. 4,000 copies printed. (Series: Ita: Fizichnyy sbirnyk, vyp. 3/8/)

Additional Sponsoring Agency: Akademiya nauk SSSR. Komissiya po spektroskopii. Ed.: Jazer, S.L.; Tech. Ed.: Saranyuk, T.V.; Editorial Board: Landsterg, G.S., Academician (Resp. Ed., Deceased), Neporent, B.S., Doctor of Physical and Mathematical Sciences, Fabelinskiy, I.L., Doctor of Physical and Mathematical Sciences, Fabelinskiy, V.A., Doctor of Physical and Mathematical Sciences, Kornitskiy, V.G., Candidate of Technical Sciences, Rayskiy, S.M., Candidate of Physical and Mathematical Sciences, Klimovskiy, L.K., Candidate of Physical and Mathematical Sciences, Miliyanovich, V.S., Candidate of Physical and Mathematical Sciences, and Glauberman, A. Ye., Candidate of Physical and Mathematical Sciences.

Card 1/30

Fentin, Yu. A., V.M. Tatevskiy, and B.A. Pordyshev. Study of Rotational Isomerism by Means of Spectroscopy	300
Kovalev, I.P. Vibrational Spectra and Potential Energy Constants of Monosilane and Its Deutero-derivatives	304
Veyts, I.V., and L.V. Gurych. Energy Dissociation and Basic Electron States of Alkali Earth Metal Oxides	305
Yakovleva, A.V., and I.I. Gromova. Nitrogen Fluorescence Under the Influence of Short-wave Radiation	308
Dianov-Klokov, V.I. Absorption Spectra of Liquid Oxygen	310
Koronkevich, V.F. Experimental Determination of Coefficients of the Dispersion Formula for Normal Air	313

"APPROVED FOR RELEASE: 03/20/2001

CIA-RDP86-00513R000617420006-5

GURVICH I V

APPROVED FOR RELEASE: 03/20/2001

CIA-RDP86-00513R000617420006-5"

"APPROVED FOR RELEASE: 03/20/2001

CIA-RDP86-00513R000617420006-5


GURVICH, L.V.

APPROVED FOR RELEASE: 03/20/2001

CIA-RDP86-00513R000617420006-5"

"APPROVED FOR RELEASE: 03/20/2001

CIA-RDP86-00513R000617420006-5



APPROVED FOR RELEASE: 03/20/2001

CIA-RDP86-00513R000617420006-5"

SOV/137-58-8-16357

Translation from: Referativnyy zhurnal, Metallurgiya, 1958, Nr 8, p 17 (USSR)

AUTHORS: Veyts, I.V., Gurvich, L.V.

TITLE: On the Problem of the Energy of Dissociation of the Basic Electronic States of the Oxides of Alkaline-earth Metals (K voprosu ob energii dissotsiatsii i osnovnykh elektronnykh sostoyaniyakh okislov shchelochnozemel'nykh metallov)

PERIODICAL: Fiz. sb. L'vovsk. un-t, 1957, Nr 3 (8), pp 305-308

ABSTRACT: The energy of dissociation of the molecules of MgO, CaO, SrO, and BaO was determined by the measurement of the constants of the equilibrium of the dissociation reactions of these oxides in the flame of H₂ with O₂ and air, as well as of acetylene with O₂ and with air. The relationships of log K_p to 1/T for the reaction of dissociation of oxides were plotted. The values for the energy of dissociation, established through the log K_p-1/T relationship in a broad temperature range, have no relation to the basic state of the oxides of the metals examined. 1. Alkaline earth metal oxides--Ionization 2. Alkaline earth metal oxides--Energy 3. Flames--Ionizing effects Yu.L.

Card 1/1

Gurvich, L. V.

76-10-20/34

AUTHORS: Veyts, I. V., Gurvich, L. V.

TITLE: On the Dissociation Energies of Oxide Molecules of the Alkaline Earth Elements (K voprosu ob energiyakh dissotsiatsii molekul okislov shchelochno-zemel'nykh elementov).

PERIODICAL: Zhurnal Fizicheskoy Khimii, 1957, Vol. 31, Nr 10, pp. 2306-2311 (USSR)

ABSTRACT: The dissociation energies of CaO and SrO were investigated here on the basis of measurements of the equilibrium constants of their dissociation in a $\text{CO}+\text{O}_2$ -flame in which the partial pressure of the hydroxyl is considerably lower than in other flames with a temperature of $\sim 3000^\circ\text{K}$. The values $D_0(\text{CaO})$ and $D_0(\text{SrO})$ obtained by the investigations in flames with a different hydroxyl content are compared. It is shown that the presence of the CaOH- and SrOH-molecules does not influence the dissociation energy values to be determined of the calcium- and strontium oxides. Furthermore it is shown that the gas temperature in the exterior flame cone of $\text{CO}+\text{O}_2$ which was measured according to the Ornstein

CARD 1/2

On the Dissociation Energies of Oxide Molecules of the Alkaline Earth Elements 76-10-20/34

method agrees well with the theoretically computed equilibrium temperature. On the strength of an analysis of the data concerning the dissociation energies of molecules of the oxides of the alkaline earth elements which were obtained according to different methods it is shown that the most precise values were obtained on the strength of the investigation of the dissociation equilibrium of the corresponding oxides in the flames. There are 4 tables and 3 Slavic references.

ASSOCIATION: Institute for Mineral Fuels
Moscow (Institut goryuchikh iskopayemykh,
Moskva).

SUBMITTED: July 27, 1956

AVAILABLE: Library of Congress

CARD 2/2

GURVICH, L. V.

AUTHORS: Gurvich, L. V., Veyts, I. V.

20-5-25/48

TITLE: Spectroscopic Investigation of the $\text{NaCl} \rightleftharpoons \text{Na} + \text{Cl}$
Reaction Equilibrium in the $\text{H}_2 + \text{Cl}_2$ Flame and the Dissociation
Energy of NaCl (Spektroskopicheskoye issledovaniye ravnovesiya
reaktsii $\text{NaCl} \rightleftharpoons \text{Na} + \text{Cl}$ v plameni $\text{H}_2 + \text{Cl}_2$ i energiya
dissotsiatsii NaCl).

PERIODICAL: Doklady AN SSSR, 1957, Vol. 116, Nr 5, pp. 811-812 (USSR)

ABSTRACT: First the results of several respective works are referred
to. The investigation was carried out in a flame fed with
the fuel compound $1,00 \text{ H}_2 + 0,80 \text{ Cl}_2 + 0,016 \text{ H}_2\text{O}$ (liquid).
The temperature of this flame was 2450° K . The partial
pressure of atomic chlorine was $3,37 \cdot 10^{-2}$ atm.
The determination of the flame temperature from the reaction
of the D-bands of Na at a distance of from 3-6 mm above
the reaction zone supplied the values $2350 - 2550^\circ \text{ K}$. For
the investigation of the reaction equilibrium $\text{NaCl} \rightleftharpoons \text{Na}$
+ Cl diluted solutions of two sodium salts ($1,02 \cdot 10^{-3}$ N-

Card 1/4

Spectroscopic Investigation of $\text{NaCl} \rightleftharpoons \text{Na} + \text{Cl}$ 20-5-25/48
Reaction Equilibrium in the $\text{H}_2 + \text{Cl}_2$ Flame and the Dissociation
Energy of NaCl.

solution of NaCl and $1,19 \cdot 10^{-3}$ N-solution of Na_2CO_3) were introduced to the flame. The partial pressure of Na in the flame was determined from the absolute intensity of the resonance bands 5890 and 5896 Å in the flame spectrum some mm above the reaction zone. The methods used for measurements and the treating of experimental data was already discussed in a preliminary work (reference 1). The values P_{Na} of the partial pressure found this way are listed in a table. The same table contains the values of $P_{\Sigma \text{Na}}$ - the sum of the partial pressures of Na and its compounds in the flames of the gases. The values of $P_{\Sigma \text{Na}}$ are calculated from the data of the sodium salt introduced to the flame, from the composition of the compound feeding the flame and from the composition of the flame gases. Using the values of P_{Na} and the partial pressures of the gaseous components of the combustion products of the hydrochloric acid flame one can

Card 2/4

Spectroscopic Investigation of $\text{NaCl} \rightleftharpoons \text{Na} + \text{Cl}$ 20-5-25/48

Reaction Equilibrium in the $\text{H}_2 + \text{Cl}_2$ Flame and the Dissociation
Energy of NaCl.

show that Na in the case of equilibrium in the flame is present only in form of NaCl, Na^+ and Na and that the partial pressures of NaH, NaOH and Na_2Cl_2 are so small that they can be neglected. A further table contains the values of P_{Na^+} , $P_{\text{NaCl}} = P_{\Sigma \text{Na}} - P_{\text{Na}^+} - P_{\text{Na}}$, the constant of the equilibrium of reaction $\text{NaCl} \rightleftharpoons \text{Na} + \text{Cl}$ with the flame temperature as well as of $D_0(\text{NaCl}) = T(\Phi_{\text{Na}}^* + \Phi_{\text{Cl}}^* - \Phi_{\text{NaCl}}^* - R \ln K_p)$. The comparison of the results found in this work with one another as well as with the results obtained by means of other methods speaks in favor of the following: In the outer core of the flame conditions must be reached which are close to equilibrium, and the dissociation energies of the metal chlorides can be investigated in a hydrochloric acid flame. There are 1

Card 3/4

Spectroscopic Investigation of $\text{NaCl} \rightleftharpoons \text{Na} + \text{Cl}$ 20-5-25-48
Reaction Equilibrium in the $\text{H}_2 + \text{Cl}_2$ Flame and the Dissociation
Energy of NaCl.

table, and 7 references, 6 of which are Slavic.

PRESENTED: May 4, 1957, by V. N. Kondrat'yev, Academician

SUBMITTED: April 25, 1957.

AVAILABLE: Library of Congress

Card 4/4

GURVICH, L.V.

AUTHORS: Matveyev, M.A., Rabukhin, A.I., Gurvich, L.V. 72-2-5/20

TITLE: Ceramic Lining of Vibration Mills (Keramicheskaya futerovka vibromel'nits).

PERIODICAL: Steklo i Keramika, 1958. Nr 2, pp. 10-13 (USSR)

ABSTRACT: In order to produce a vibration mill that is proof against wear, and also in order to avoid the metal- or rubber lining in vibration mills such as are in use now, a method of fastening a ceramic lining had to be found. For test purposes the vibration mill M - 200 - 1.5 with a separate vibrator was developed by SKB VNIITISM for lining with plates made of various materials (see total view fig. 1). The lining plates are shown in form of drawings in figs. 2, 3, 4 and 5. For the fastening of these plates various kinds of adhesives were tested, and it was found that an adhesive based upon resin ED - 6 gave the best results. The production of this adhesive is then described in detail, as also, in fig. 6, the manner of fastening the ceramic plates. Tests were then carried out with lining plates of different origins. Fig. 7 shows holders with glued-on uralite plates. Figs. 8, 9 and 10 show linings of uralite, porcelain, and earthenware after having been in operation

Card 1/2

Ceramic Lining of Vibration Mills

72-2-5/20

for test purposes, without interruption, for 110 hours, without any damage having been found. There are 10 figures and 1 Slavic reference.

ASSOCIATION: VNIITISM

AVAILABLE: Library of Congress

Card 2/2

AUTHOR: Gurvich, L.V.

SOV/51-5-2-18/26

TITLE: The Absolute Probabilities of Transitions in an Atom of Tl
(Ob absolyutnykh veroyatnostyakh perekhodov atoma Tl)

PERIODICAL: Optika i Spektroskopiya, 1958, Vol 5, Nr 2, pp 205-207 (USSR)

ABSTRACT: The absolute values of the oscillator strengths of the 3776 and 5350 Å lines of Tl had been earlier determined experimentally (Refs 1-5) and by calculation (Refs 4, 6). Table 1 gives the values obtained by the authors of Refs 1-5. The most reliable set of experimental results is that of Kvater (Ref 3) obtained by the method of "hooks". Comparison of Kvater's f values with those calculated theoretically, as well as with those found experimentally for Ga and In (Ref 7), shows them to be too small. Kvater himself pointed out the inexactness of determination of the absolute f values because of lack of reliable data on the saturated vapour pressure of Tl. The author shows that, employing simple thermodynamical relationships, it is possible to determine the absolute values of f by the method of "hooks" without using any data on the vapour pressures of metals. It is sufficient to

Card 1/3

The Absolute Probabilities of Transitions in an Atom of Tl SOV/51-5-2-18/26

measure the values of Nf in a wide range of temperatures and to know the thermodynamical properties of the metal in solid and liquid states. The author describes this thermodynamic method of treatment of experimental data and applies it to find the absolute values of f for Tl using Kvater's experimental results (Ref 3). Table 2 lists the values of the absolute oscillator strengths for the 3776 and 5350 lines of Tl at temperatures of 885-1336°K (columns 4 and 6 of Table 2) obtained in this way. The mean value of f for the 3776 line was 0.125 and for the 5350 line it was 0.135. The latter value agrees well with the results of theoretical calculations as well as with the value found by Stephenson (Ref 4). The values of the oscillator strengths for the 3776 and 5350 Å lines of Tl make it possible to correct the absolute values of the oscillator strengths for other Tl lines. The described thermodynamical method for calculation of f was applied to the experimental results of Ostrevskiy and Penkin (Ref 7) for In. The values thus obtained differ by only 10% from those

Card 2/3

SOV/51-5-2-18/26

The Absolute Probabilities of Transitions in an Atom of Tl

given by the latter authors in Ref 7, which confirms that in the case of In reliable values of the saturated vapour pressure were available. There are 2 tables and 11 references, 7 of which are Soviet, 1 Danish, 1 English, 1 American and 1 Swiss.

ASSOCIATION: Institut goryuchikh iskopayemykh, AN SSSR (Institute of Fuel Minerals, Academy of Sciences of the U.S.S.R.)

SUBMITTED: February 6, 1958

Card 3/3 1. Atoms--Properties 2. Thallium--Properties 3. Perturbation theory
--Mathematical analysis

AUTHORS: Gurrich, L. V., Veyts, I. V. SOV/48-22-6-9/28

TITLE: The Determination of the Dissociation Energies of Molecules by the Investigation of the Equilibrium of Their Dissociation in Flames (Opradeleniye energiy dissotsiatsii molekul na osnovanii izucheniya ravnovesiya ikh dissotsiatsii v plamenakh)

PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya fizicheskaya, 1958, Vol. 22, Nr 6, pp. 673-676 (USSR)

ABSTRACT: When studying earlier papers (Refs 1-6) we find that one of the best methods of determining dissociation energies is based upon determination of the partial pressure P_M of the metal according to the intensity of its lines in the flame spectrum. Basing on the assumption that in flames metal can only exist in form of the atoms M and the molecules MX, partial pressure P_{MX} of the compound under investigation is determined according to the following formula: $P_{MX} = P_M - P_M$, where P_M denotes the general pressure of the metal compounds in the flames. It is pointed out (Refs 5-7) that partial pressure can be determined according to the known equilibrium constant of the dissociation reactions of the compounds.

Card 1/3

The Determination of the Dissociation Energies of
Molecules by the Investigation of the Equilibrium
of Their Dissociation in Flames

SOV/48-22-6-9/28

On the basis of the examples of calculation given it is found that the dissociation energy MX can be determined by the method mentioned only if $D_0(MX) \geq T(\Delta \phi_T + 1.38 - R \ln PX)$. The presence of other metal compounds impairs the accuracy of this method (Ref 8). The method was employed for the investigation of dissociation energies of the diatomic oxides of the elements of group 3 (BO, AlO, GaO, InO, TlO) in the case of flame compositions of $C_2H_2+O_2$, H_2+O_2 , $CO+O_2$. Measurements were carried out as described (Refs 5,6). The metals were introduced into the flames as solutions of their sulfuric acid- or chromium salts and boron in form of $Na_2B_4O_7$. The partial pressure of metals in flames was determined on the basis of measurements carried out of the intensity of resonance lines with the transitions $^2S_{1/2} \rightarrow ^2P_{1/2}, 3/2$. Results are shown by tables. In conclusion it is pointed out that this paper confirms the possibility of accurately determining dissociation energies of molecules in flames.

Card 2/3

The Determination of the Dissociation Energies of
Molecules by the Investigation of the Equilibrium
of Their Dissociation in Flames

SOV/48-22-6-9/28

At the same time it is mentioned that by comparing stable molecules BO , AlO , GaO and InO it can be proved that the theory developed by H. G. Howell (Ref 20) concerning the low dissociation energy of InO is wrong. There are 24 references, 9 of which are Soviet.

ASSOCIATION: Institut goryuchikh iskopayemykh Akademii nauk SSSR (Institute of Mineral Fuels, AS USSR)

1. Metals--Ionization
2. Energy--Measurement
3. Metals--Spectra
4. Flames--Spectra
5. Mathematics

Card 3/3

SOV/76-32-11-11/32

5(4)

AUTHORS:

Veyts, I. V., Gurvich, L. V., Rtishcheva, N. P.

TITLE:

Thermodynamic Properties of Magnesium, Calcium, Strontium, Barium and of Their Oxides and Monohydrides in Gaseous State (Termodinamicheskiye svoystva magniya, kal'tsiya, strontsiya, bariya, ikh okislov i monogidridov v gazoobraznom sostoyanii)

PERIODICAL:

Zhurnal fizicheskoy khimii, 1958, Vol 32, Nr 11, pp 2532-2542 (USSR)

ABSTRACT:

The thermodynamic properties of the substances mentioned in the title were determined according to the methods of static thermodynamics at 293.16; 298.16; 400 (100) and 3500°K. From the data obtained the equilibrium constants of the dissociation of the corresponding diatomic gases in the investigated temperature range were calculated. The values for the

thermodynamic potential $\phi_T^* = - \frac{Z_T^* - H_O^*}{T}$, the entropy S_T^* and the change of the enthalpy $H_T^* - H_O^*$ were determined as well.

The determination of the thermodynamic functions of the diatomic gases was carried out according to the table method by

Card 1/2

SOV, 76-32-11-11/32
Thermodynamic Properties of Magnesium, Calcium, Strontium, Barium and of
Their Oxides and Monohydrides in Gaseous State

Gordon and Barnes (Ref 25). In the calculations the values of the ϕ^* potentials for atomic oxygen and hydrogen recommended by the Byuro standartov SSHA (Bureau of Standards USA) were used. The authors thank I. G. Baybuz and V. S. Shmeleva. There are 11 tables and 29 references, 5 of which are Soviet.

ASSOCIATION: Institut goryuchikh iskopayemykh, Moskva (Institute of
Mineral Fuel, Moscow)

SUBMITTED: May 4, 1957

Card 2/2

BROUNSHTEYN, B.I.; GURVICH, L.V.; YUNGMAN, V.S.; YURKOV, G.N.

Statistical methods of computing the thermodynamic functions of ideal gases. Report No. 1: General relationships of statistical thermodynamics for ideal gas. Trudy GIPKH no.42:3-10 '59. (MIRA 13:10)
(Thermodynamics) (Gases)

BROUNSHTEYN, B.I.; GURVICH, L.V.; YUNGMAN, V.S.; YURKOV, G.N.

Statistical methods of computing the thermodynamic functions of ideal gases. Report No. 2: Expression for the statistical sum based on the states of diatomic molecules. Method of direct summation based on the levels of diatomic molecules. Trudy GIPKH no.42:11-20 '59.

(MIRA 13:10)

(Gases)

(Thermodynamics)

BROUNSHTEYN, B.I.; GURVICH, L.V.; YUNGMAN, V.S.; YURKOV, G.N.

Statistical methods of computing the thermodynamic functions of ideal gases. Report 3: Approximate methods of calculating the statistical sum from the rotational states of diatomic molecules. Trudy GIPKH no.42:21-50 '59. (MIRA 13:10)

(Thermodynamics)

(Gases)

24(7), 5(2)

SOV/51-1-18/27

AUTHORS: Gurvich, L.V. and Novikov, M.M.

TITLE: On the Valence Angle of Oxygen in the HOCl Molecule (O valentnom ugle kisloroda v molekule HOCl)

PERIODICAL: Optika i spektroskopiya, 1959, Vol 7, Nr 1, pp 116-117 (USSR)

ABSTRACT: Hedberg and Badger (Ref 1) used the infrared spectrum of gaseous HOCl to deduce that the oxygen valence angle \angle HOCl is equal to 113° . The present authors are of the opinion that this angle should not be greater than 110° and they repeat Hedberg and Badger's calculations showing that best agreement with the empirical data is obtained with \angle HOCl = $104 \pm 3^\circ$. There is 1 English reference.

SUBMITTED: November 5, 1958

Card 1/1

24814

S/081/61/000/011/005/040

B105/B203

245300

AUTHORS: Yungman, V. S., Gurvich, L. V., Kvlividze, V. A.,
Rtishcheva, N. P.

TITLE: Thermodynamic functions of monoatomic and diatomic gases
in a wide temperature range. IV. N^+ , N_2^+ , and NO^+ in
ideal state up to 20,000° K

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 11, 1961, 40, abstract
115 292 (Sb. tr. Gos. in-ta prikl. khimii, 1960, vyp. 46,
15 - 28)

TEXT: The thermodynamic functions (Φ_T^* , S_T^0 , and $H_T^0 - H_0^0$) of N^+ , N_2^+
and NO^+ in ideal state up to 20,000° K at a pressure of 1 atm were
calculated on an electronic computer. In the values Φ_T^* and S_T^0 at
 $T \leq 10,000^\circ K$, the error does not exceed 0.01 cal/mole/deg, and at
 $T = 20,000^\circ K$ it does not exceed 0.2 cal/mole/deg. The values of the
logarithms of the equilibrium constants for the ionization of N, N_2 , NO
and the dissociation of N_2^+ and NO^+ are given. [Abstracter's note:
Card 1/1]

83691

S/076/60/034/008/003/014
B015/B054

5.4700 (1273)

AUTHOR: Gurvich, L. V. (Moscow)

TITLE: Determination of the Sublimation Heats of Metals on the Basis of Measurements of Anomalous Dispersion by the Method of Rozhdestvenskiy. Thermodynamic Properties and Sublimation Heats of Gallium, Indium, and Thallium

PERIODICAL: Zhurnal fizicheskoy khimii, 1960, Vol. 34, No. 8, pp. 1691-1698

TEXT: It is shown that the method developed by D. S. Rozhdestvenskiy (Ref. 3) can be used to determine the sublimation heats of Ga, In, and Tl. The method is based on the determination of the product $N \cdot f$ (N = number of atoms of the metal vapor in 1 cm^3 , f = absolute value of the forces of the oscillator for the spectral line of the atom in which the interference peaks are measured) from the measured values of the distance between the interference peaks. The probability of optical transitions of atoms and the forces of oscillators are constant for any

Card 1/3

83691

Determination of the Sublimation Heats of Metals on the Basis of Measurements of Anomalous Dispersion by the Method of Rozhdestvenskiy. Thermodynamic Properties and Sublimation Heats of Gallium, Indium, and Thallium

S/076/60/034/008/003/014
B015/B054

spectral line, and it is possible to write down $\ln NfT = A - \frac{B}{T}$ (1)

(R = gas constant, A and B = coefficients, T = temperature), where B = heat of vaporization of the metal. Thus, the sublimation heat of the metal for 0°K can be determined with high accuracy if the thermodynamic properties of the metal for the solid and gaseous state are known. The data of the oscillator forces of the Ga, In, and Tl atoms were taken from publications, and the thermodynamic properties in the gaseous state were calculated statistically (Table 1, data calculated for 293.15°, 298.16°, 400° to 3500°K), as well as for the condensed state (Table 2, values for 298.16° to 1500°K). The values for N-f of the resonance lines of Ga, In, and Tl were investigated by Yu. I. Ostrovskiy and N. P. Penkin, as well as G. S. Kvater, at the Leningradskiy universitet (Leningrad University). With the aid of these data, the following values were calculated.

Card 2/3

83691

S/076/60/034/008/003/014
B015/B054

Determination of the Sublimation Heats of Metals on the Basis of Measurements of Anomalous Dispersion by the Method of Rozhdestvenskiy. Thermodynamic Properties and Sublimation Heats of Gallium, Indium, and Thallium

ΔH_0° (Ga, gas) = 59000 \pm 1000 cal/gram atom; ΔH_0° (In, gas) = 57700 \pm 600 cal/gram atom; ΔH_0° (Tl, gas) = 43150 \pm 250 cal/gram atom. The author mentions a paper by L. P. Lyubimov and Yu. N. Lyubotov (Ref. 25) which was published after the present paper had been printed. There are 2 tables and 27 references: 6 Soviet, 10 US, 1 British, 8 German, and 1 Belgian. ✓

ASSOCIATION: Akademiya nauk SSSR, Institut goryuchikh iskopayemykh
(Academy of Sciences of the USSR, Institute of Mineral
Fuels)

SUBMITTED: September 12, 1958

Card 3/3

32326

S/081/61/000/024/008/086

B138/B102

11.5100
11.3000
AUTHORS:

~~Gurvich, L. V.~~, Yungman, V. S., Prozorovskiy, Ye. A.,
Vorob'yev, B. A.

TITLE:

Calculation of the thermodynamic functions of diatomic gases
at elevated temperatures by direct summation on an electro-
nic machine

PERIODICAL:

Referativnyy zhurnal. Khimiya, no. 24, 1961, 62, abstract
24B422 (Tr. In-ta goryuchikh iskopayemykh AN SSSR, v. 12,
1961, 196 - 205)

TEXT: A very rapid and precise method is proposed for the calculation
of the thermodynamic function tables of diatomic perfect gases at tempera-
tures of up to 20,000 to 25,000°K. The statistical sums are calculated,
for the rotational vibrational and electron states of the molecule in
question, by direct summation through the really existant energy levels,
using a high-speed electronic computer. For this kind of calculation the
molecular constant which most precisely describes all the energy levels of

Card 1/3

Calculation of the thermodynamic ...

32326
S/081/61/000/024/008/086
B138/B102

the molecule must be known, as also the highest values of the quantum numbers up to which summation is to be made. A method is described for calculating vibrational constants and maximum vibrational quantum numbers $v(\max)$ using the conditions for the convergence of the vibrational levels toward the dissociation limit. A method has been developed for calculating values of rotational quantum numbers $J(\max)$ for each vibrational state, using the properties of the effective potential curves of the rotating molecule. As an example some results are given of the calculation of the main state $x^3\Sigma_g^-$ of an O_2 molecule. In particular, to describe the energy of vibrational levels (in cm^{-1}) the equation $G_o(v) = 1568.077 v - 11.706 v^2 - 0.00255 v^3 + 0.00224 v^4 - 0.0000821 v^5$ is derived, which converges towards the 41261 cm^{-1} limit at $v(\max) = 42$ (experimental values of dissociation energy of O_2 are $41260 \pm 15 \text{ cm}^{-1}$). $J(\max)$ values are found for all v . The thermodynamic functions of molecular oxygen are given for the following temperatures: 5000°K (63.395 and 73.038), 10000°K (70.457 and 79.942).

Card 2/3

32326

Calculation of the thermodynamic ...

S/081/61/000/024/008/086
B138/B102

15000°K (74.229 and 83.255) and 20000°K (76.746 and 83.203) (values in brackets are for the isobaric-isothermal potential ϕ_T^* and entropy S_T^0 respectively, in cal/mol, degree. [Abstracter's note: Complete translation.] ✓

Card 3/3

GURVICH, L.V. (Moskva); KVLIVIDZE, V.A. (Moskva)

Thermodynamic functions of mono- and diatomic gases over a wide temperature range. Part 1: Method of calculating the thermodynamic functions of monoatomic gases in the ideal state. Zhur.fiz.khim. 35 no.8:1672-1680 Ag '61.
(MIRA 14:8)

(Gas dynamics)

GURVICH, L.V.; YUNGMAN, V.S. (Moskva)

Thermodynamic functions of mono- and diatomic gases in a wide range of temperatures. Part 2: Method for computing the thermodynamic functions of diatomic gases in the ideal state. Zhur.fiz.khim. 35 no.9:1927-1934 '61. (MIRA 14:10)

1. Institut goryuchikh iskopayemykh.
(Gas dynamics)

YUNGMAN, V.S.; GURVICH, L.V.; KVLIVIDZE, V.A.; PROZOROVSKIY, Ye.A.;
RTISHCHEVA, N.P. (Moscow)

Thermodynamic functions of mono- and diatomic gases in a wide
temperature range. Part 3: N, N₂ and NO in the ideal state up
to 20000 K. Zhur.fiz.khim. 35 no.10:2182-2189 0 '61.

(MIRA 14:11)

1. Akademiya nauk SSSR, Institut goryuchikh iskopayemykh.
(Nitrogen) (Nitrogen oxide) (Gas dynamics)

VEDENEYEV, Vladimir Ivanovich; GURVICH, Lev Veniaminovich; KONDRAT'YEV, Viktor Nikolayevich, akademik; MEDVEDEV, Vadim Andreyevich; FRANKEVICH, Yevgeniy Leonidovich; DRAGUNOV, E.S., red.; RYLINA, Yu.V., tekhn. red.

[Energies of chemical bond breaking. Ionization potentials and electron affinity] Energii razryva khimicheskikh svyazei. Potentsialy ionizatsii i sredstvo k elektronu; spravochnik. [By] V.I. Vedeneyev i dr. Moskva, Izd-vo Akad. nauk SSSR, 1962. 215 p.
(MIRA 16:2)

(Chemical bonds) (Ionization) (Chemical affinity)

PHASE I BOOK EXPLOITATION

SOV/6260

Gurvich, Lev Veniaminovich, Georgiy Akopovich Khachkuruzov, Vadim Andreyevich Medvedev, Inessa Veniaminovna Veyts, Georgiy Andreyevich Bergman, Vladimir Stepanovich Yungman, Nina Petrovna Rtishcheva, Lidiya Fedorovna Kuratova, Georgiy Nikolayevich Yurkov, Amaliya Abramovna Kane, Boris Fedorovich Yudin, Boris Isidorovich Brounshteyn, Viktor Feodosyevich Baybuz, Valeriy Aleksandrovich Kvlividze, Yevgeniy Aleksandrovich Prozorovskiy, and Boris Aleksandrovich Vorob'yev.

Termodinamicheskiye svoystva individual'nykh veshchestv; spravochnik v dvukh tomakh. tom 1: Vychisleniye termodinamicheskikh svoystv; tom 2: Tablitsy termodinamicheskikh svoystv (Thermodynamic Properties of Individual Substances; Reference Book in Two Volumes. v. 1: Calculation of Thermodynamic Properties; v. 2: Tables of Thermodynamic Properties). 2d ed., rev. and enl. Moscow, Izd-vo AN SSSR, 1962. 1161 and 916 p. 4000 copies printed.

Sponsoring Agencies: Akademiya nauk SSSR. Institut goryuchikh iskopayemykh; and Gosudarstvennyy komitet Soveta Ministrov SSSR

Card 1/93

Thermodynamic Properties (Cont.)

SOV/6260

po khimii. Institut prikladnoy khimii.

Resp. Ed.: V. P. Glushko, Academician, L. V. Gurvich, G. A. Khachkuruzov, I. V. Veyts, and V. A. Medvedev; Ed. of Publishing House: K. P. Gurov; Tech. Ed.: V. G. Laut.

PURPOSE: This reference book may be used in scientific-research and experimental-design work in institutes, design offices, and schools of higher education, as well as for training specialists in chemical thermodynamics and thermal physics.

COVERAGE: Volume 1 of this work deals with methods for calculating thermodynamic properties and with the selection of constants required for the calculations. Volume 2 contains tables of thermodynamic properties (reduced thermodynamic potential, entropy, enthalpy, and the logarithm of the dissociation or ionization constants of equilibrium) compiled, where data were lacking, on the basis of published and unpublished material from a number of Soviet research institutes. Thermodynamic properties for the ideal gas

Card 2/9₃

Thermodynamic Properties (Cont.)

SOV/6260

state are presented in table form for 335 gases, 44 liquids, and 45 solids compounded from 33 chemical elements and their isotopes, viz.: H, D, T, He, Li, Be, B, C, N, O, F, Ne, Na, Mg, Al, Si, P, S, Cl, Ar, K, Ca, Br, Kr, Re, Sr, Zr, I, Xe, Cs, Ba, Hg, and Pb. Thermodynamic properties are given for the following 22 gases in the range from room temperature to 20,000°K: H, H⁺, H⁻, O, O⁺, H₂, O₂, OH, OH⁺, H₂O, N, N⁺, N₂, N₂⁺, NO, NO⁺, C, C⁺, CO, CO⁺, and e⁻; for the 14 least stable gases up to 4000°K; and for the remaining 299 gases up to 6000°K. Virial coefficients for 34 gases are also given up to 6000°K.

TABLE OF CONTENTS (Volume 1) [Abridged]:

Foreword	11
Introduction	19
PART I. METHODS OF CALCULATING THE THERMODYNAMIC PROPERTIES OF INDIVIDUAL SUBSTANCES	

Card 3/93

YUNGMAN, V.S.; GURVICH, I.V.; STRECHENVA, L.P.

Thermodynamic properties of gaseous compounds of nitrogen with
hydrogen (NH, NH₂, and N₂H₄). Trudy GIKH no.49:20-37 '62.
(MIRA 17:11)

ONITSKI, I.V.; K. S. LITVIN, A.A.; KULIKOV, I.V.; ... LITVIN, A.A.; LITVIN, A.A.,
P.P.; LITVIN, V.S.

Thermodynamic functions of mono- and diatomic gases within a wide
range of temperatures. Part 6: O, O^+, O_2 , and O_2^+ in the ideal state
up to $20\,000^\circ\text{K}$. Trudy GIFKH no.49:38-60 '62.

(MIRA 17:11)

Thermodynamic functions of mono- and diatomic gases within

a wide range of temperatures. Part 7: C, C⁺, CO, CO⁺ in the
ideal state up to 20 000⁰ K. Trudy GIPKH no.49:61-83 '62.
(MIRA 17:11)

32643
S/076/62/036/001/015/017
B119/B101

11.4100

AUTHORS: Gurvich, L. V., Kvlividze, V. A., and Rtishcheva, N. P.

TITLE: Thermodynamic functions of monatomic and diatomic gases within a wide temperature interval. V. Alkali metals in the state of an ideal gas up to 10,000°K

PERIODICAL: Zhurnal fizicheskoy khimii, v. 36, no. 1, 1962, 219 - 222

TEXT: The thermodynamic functions of gaseous Li, Na, K, Rb, and Cs (Φ_T^* , S_T^0 , $H_T^0 - H_O^0$) were calculated for temperatures of up to 10,000°K by the method of L. V. Gurvich and V. A. Kvlividze (Zh. fiz. khimii, 35, 1672, 1961). When calculating the statistical sums and the figures derived from them, allowance was made for all the electron states of the individual gases, which are related to the transition of the valence electron into a state with the principal quantum number $n \leq n_{\max}$. $n_{\max} = 2.461 \cdot T^{1/6}$. ✓

$\log K_p$ was calculated from $\log K_p = 0.21854 [\Phi_T^*(M^+) + \Phi_T^*(e) - \Phi_T^*(M) - \frac{I}{T}]$, where $\Phi_T^*(M)$, $\Phi_T^*(M^+)$, and $\Phi_T^*(e)$ are the values of Φ_T^* of the gaseous alkali metal,

Card 1/2

DECLASSIFIED: APRIL 15, 1965

Card 2/2

L 8550-65 EWT(1)/EWT(m) ASD(a)-F/AFTC(b)/AFWL/AFMD(p)/AFETR/SSD/EBD(dp)/
ESD(t)/RAEM(t) JD/JW
ACCESSION NR: AR4044045 S/0098/63/000/011/E002/1002

SOURCE: Ref. zh. Fizika, Abs. 11E6

AUTHOR: Gurvich, L. V.; Vorob'yev, B. A.; Kvilidze, V. A.; Prozorovskiy, Yu. A.;
Rtishcheva, N. P.; Yunguan, V. S.

TITLE: Thermodynamic functions of monatomic and diatomic gases in a broad
temperature range. VI. O, O+, and O₂ in the ideal state to 20,000°K

CITED SOURCE: (Sb. tr.) Gos. in-ta prikl. khimii, v. 49, 1962, 38-60

TOPIC TAGS: thermodynamic function, monatomic gas, diatomic gas, high speed
computer, computer

TRANSLATION: Gives the results of calculations of thermodynamic functions (Ω_T , S_T^0
and $H_T^0 - H_0^0$) O, O+, O₂, and O₂+, made, in accordance with previously-described methods
(Journal of Abstracts, Physica, 1D4; 2D10), on a high-speed computer. Bibliography:
67 references.

SUB CODE: GO, TD

ENCL: 00

Card 1/1

L 8551-65 EWT(1)/EWT(m) SSD/AFMD(p) AFWL/AFETR/ASD(a)-5/AFTC(b)/ESD(dp)/
ESD(t)/RAEM(t) JD/JW
ACCESSION NR: AR4044046 S/0058/63/000/011/E002/E002

SOURCE: Ref. zh. Fizika, Abs. HE7

AUTHOR: Gurvich, L. V.; Kvilidze, V. A.; Prozorovskiy, Ye. A.;
Rtishcheva, N. P.

TITLE: Thermodynamic functions of monatomic and diatomic gases in a broad
range of temperatures. VII. C, C⁺, CO, CO⁺ in the ideal state to 20,000°K

CITED SOURCE: (Sb. tr.) Gos. in-ta prikl. khimii, vyip. 49, 1962, 61-83

TOPIC TAGS: thermodynamic function, monatomic gas, diatomic gas, computer

TRANSLATION: Gives the results of calculations of the thermodynamic functions
(Φ_r° , S_r° and H_r°) C, C⁺, CO and CO⁺ made on an electronic computer in accordance
with previously described methods (Journal of Abstracts, Physics, 1962, 1D4;
2D10). Bibliography: 73 references. Part VI: see abstract 11136.

SUB CODE: GC, TD
Card 1/1

ENCL: 00

S/051/63/014/002/023/026
E039/E120

AUTHORS: Gurvich, L.V., and Shenyavskaya, Ye.A.

TITLE: The electron spectrum of scandium monofluoride

PERIODICAL: Optika i spektroskopiya, v.14, no.2, 1963, 307-308

TEXT: This investigation was carried out in order to provide information on the spectra of diatomic compounds of elements of subgroup IIIb with halogens. A discharge tube containing ScF_3 and metallic Sc, with He and A as a discharge carrier, was used as a light source. Spectra were obtained using an MCN-28 (ISP-28) spectrograph and the optimum conditions were: cathode (Armco iron) 6 mm diameter, 30 mm long, 410 V, 350 mA, He at a pressure of 6 mm Hg. In the region of 2850 Å a group of bands was obtained which had not previously been observed. Their intensity was too low for analysis and they were overlapped by iron lines. More satisfactory results were obtained using a quartz tube with a 10 mm diameter capillary 150 mm long and heated externally by a nichrome helix. The tube contained a mixture of Sc and ScF_3 and was sleeved with platinum in order to prevent the fluoride reacting with the quartz. Optimum conditions were: He and A at 2 mm

Card 1/2

S/051/63/014/002/023/026
E039/E120

The electron spectrum of scandium...

pressure; 1.2 - 1.3 amp at 2.4 - 3 kV. The ScF spectrum in the range 2550-3000 Å was observed in the second order using a ДФС-8 (DFS-8). Four groups of bands were observed in the violet and one band system in the red. Greatest intensity was at about 2880 Å. An analysis of the vibrational structure of the spectra was carried out. Calculated values of the molecular constants for ScF are:

$$\omega_e'' = 734.3; \quad \omega_e'' x_e'' = 3.5; \quad e = 35013.4; \quad \omega_e' = 582.6;$$

$$\omega_e' x_e' = 6.1 \text{ cm}^{-1}. \quad \text{It is shown that the energy of dissociation of}$$

ScF is ~4.5 to 6 eV.

There is 1 table.

SUBMITTED: July 30, 1962

Card 2/2

GURVICH, L.V.; YUNGMAN, V.S.

Temperature dependence of the thermodynamic functions of an ideal gas. Teplofiz. vys. temp. 2 no.1:118-119 Ja-F '64. (MIRA 17:3)

1. Nauchno-issledovatel'skiy institut vysokikh temperatur.

GUR/ICH, L.V.; RYABOVA, V.G.

Dissociation energy of the BaCl molecule. Teplofiz. vyss. temp.
2 no.2:215-218 Mr-Ap '64. (MIRA 17:6)

ACCESSION NR: AP4042467

S/0294/64/002/003/0401/0405

AUTHORS: Gurvich, L. V.; Ryabova, V. G.

TITLE: The determination of metal halide dissociation energies on the basis of equilibrium reaction studies in flames. 1. Dissociation energy of BaF

SOURCE: Teplofizika vysozikh temperatur, v. 2, no. 3, 1964, 401-405

TOPIC TAGS: equilibrium reaction, hydrogen air flame, barium, partial pressure, fluorine, dissociation rate, atomic line intensity

ABSTRACT: The equilibrium reaction rate of Ba in a hydrogen-air flame, $aH_2 + bO_2 + cN_2 + dH_2O$, with the addition of $(C_2F_5)_3N$ vapor, was investigated experimentally. Barium was added to the flame in the form of 0.02 M $BaCl_2$ solution. The absolute value of Ba partial pressures in eight different flames, at 20-mm height, was determined by the absolute line intensity $\lambda = 5535 \text{ \AA}$ with the additions of several fluorine rates (5.5, 15.8, 35, and 50 mg/min of $(C_2F_5)_3N$) and also in the absence of fluorine. From this the dissociation energy $D_0(BaF)$ of barium fluoride was determined, using

Card 1/2

ACCESSION NR: AP4042467

the expression

$$D_0(\text{BaF}) = T[\Delta\Phi_r^\circ - R \ln K_p(\text{BaF})],$$

which gave a value of 147.6 ± 1.7 kcal/mol. The dissociation energy was also determined from the ratio $I_{\text{Ba}}/I'_{\text{Ba}}$ (relative atomic line intensity) as a function of the partial pressure $P_{\Sigma F}$ which in turn gave a value of 147.6 ± 2.2 , in excellent agreement with the first method. Orig. art. has: 5 formulas, 2 tables, and 1 figure.

ASSOCIATION: Nauchno-issledovatel'skiy institut vyssokikh temperatur (Scientific Research Institute of High Temperatures)

SUBMITTED: 06Apr64

ENCL: 00

SUB CODE: FP

NO REF SOV: 006

OTHER: 008

Card

2/2

E 40786-65 EWG(j)/EWT(m)/EPT(c)/EWG(m)/EPR/T/EWP(t)/EWP(b) Pr-4/Pg-4

IJP(c)/RPL JD/JW/JG/RM

ACCESSION NR: AP4044520

S/0294/64/002/004/0540/0548

AUTHORS: Gurvich, L. V.; Ryabova, V. G.

TITLE: Investigation of dissociation energies of gallium and indium oxygen compounds in flames during equilibrium reaction I. Experimental method and investigation of $aH_2 + bO_2 + cN_2 + dH_2O$ flames

SOURCE: Teplofizika vysokikh temperatur, v. 2, no. 4, 1964, 540-548

TOPIC TAGS: combustion, flame, gallium oxide, indium oxide, dissociation, metallized flame, reaction rate constant, line spectrum, partial pressure, UM 2 monochromator, M 106/1 millivoltmeter, Orekh power source, LP 3 lamp, SI 16 lamp

ABSTRACT: The relative and absolute spectral line intensities of metal atoms in flames were studied in order to determine the dissociation energy of metallic compounds in the combustion products of the flames. It is shown that if the metal Me forms a compound MeX with a radical in the flame ($O, OH,$ or H) one can calculate the reaction rate constant K_p in $MeX \rightleftharpoons Me + X$ by determining the partial pressure of the metal. This in turn can be accomplished from the absolute spectral line intensities of the metal atoms in the flame obtained by comparing relative intensities with standard calibrated sources of continuous spectra. Thus

Card 1/3

L 40786-65

ACCESSION NR: APL044520

$$p_{\text{He}} = \frac{k\lambda}{hc g_m A_{mn}} \tau \frac{T}{T} Q(T) e^{-E_m/T} B I_{\lambda, \text{lamp}}^{\text{abs}}$$

and

$$I_{\lambda, \text{lamp}}^{\text{abs}} = \epsilon_c(T) \lambda^{-2} c_1 e^{-E_m/T} u_1$$

To measure these absolute and relative resonance lines, type $\text{aH}_2 + \text{bO}_2 + \text{cN}_2 + \text{dH}_2\text{O}$ flames were used and the metal was introduced in the flames as salts. The flame temperatures were measured by the spectral line-reversal technique and ranged between 1765C and 2159C. These results were checked by measuring D-line intensity of Na after introducing 5×10^{-3} M solution of NaCl in the flame. In order to determine radical concentrations in the flames, the equilibrium reaction of lithium with the combustion products was investigated. At 1600-2000C temperatures lithium exists as Li atoms and LiOH compounds with a known reaction constant. This in turn allows one to determine p_{H} and subsequently p_{OH} and p_{O} . The results of the measurement on the equilibrium reaction of Os and In with combustion products will be given in a further study. Orig. art. has: 11 formulas, 3 tables, and 1 figure.

ASSOCIATION: Nauchno-issledovatel'skiy institut vysokikh temperatur (Scientific Research Institute of High Temperatures)

Chem 273

L 40786-65

ACCESSION NR: AP4044520

SUBMITTED: 28Apr64

ENCL: 00

SUB CODE: FP, GC

NO REF SOV: 006

OTHER: 009

Card 3/3

598

L 24480-65
WW/JW/RH

EWT(m)/EPF(c)/EPR/ENP(j)/ENP(b)

Pc-4/Pr-4/Pa-4

RPL

JD/

ACCESSION NR: AP4047388

S/0294/64/002/005/0834/0835

AUTHORS: Ryabova, V. G.; Gurvich, L. V.

B

TITLE: Determining dissociation energies of metal halides on the basis on investigating the equilibrium of reactions in flames. 2. Dissociation energies of CaF , CaF_2 , SrF and SrF_2

SOURCE: Teplofizika vysokikh temperatur, v. 2, no. 5, 1964, 834-835

TOPIC TAGS: chromatographic analysis, dissociation energy, fluorine compound, halide

ABSTRACT: The authors present a continuation of research results in investigating dissociation energies of metal halides based upon studies of the equilibrium of reactions in flames (L. V. Gurvich and V. G. Ryabova, Teplofizika vysokikh temperatur, 2, No. 2, 215, 1964; and 2, No. 3, 401, 1964). Spectroscopic determination of equilibrium constants for the reactions forming CaF and SrF molecules was carried out by introducing strontium and calcium into hydrogen-air flames containing a small percentage of fluorine (about 0.4% of the flame gases). The flame composition was varied according to the formula $[a \text{ H}_2 + b \text{ O}_2 + c \text{ N}_2 + d \text{ H}_2\text{O}]$, where a, b, c, and d are varied to produce 8 different flame types. The authors described the experimental setup and combustion products in an earlier work

Card 1/2

L 24480-65

ACCESSION NR: APh047388

(Teplofizika vyssokikh temperatur 2, No. 4, 1964). In the reaction of the type $Me + HF \rightleftharpoons MeF + H$, the equilibrium constants were determined by spectroscopic measurement of partial pressures and by the variation of the relative intensity of the metal line in the flame spectra with and without fluorine addition for varying quantities of fluorine. Reaction equilibrium constants showed close agreement for all tests. The corresponding dissociation energies were found to be 135 ± 7 and 132 ± 7 kcal/mole for CaF and SrF respectively. CaF_2 and SrF_2 molecules were formed by the reaction type $MeF + HF \rightleftharpoons MeF_2 + H$ in an excess of fluorine. Consideration of partial pressures led to association energies of ≈ 136 kcal/mole for $(CaF - F)$ and ≈ 140 kcal/mole for $(SrF - F)$. Summing the dissociation and association energies yielded close agreement with earlier work performed in a different manner. Orig. art. has: 4 equations.

ASSOCIATION: Nauchno-issledovatel'skiy institut vyssokikh temperatur (Scientific Research Institute of High Temperatures)

SUBMITTED: 31Aug64

ENCL: 00

SUB CODE: GC

NO REF SOV: 004

OTHER: 001

Card 2/2

MEDVEDEV, V.A.; YUNGMAN, V.S.; VOROB'YEV, A.F.; GURVICH, L.V.;
BERGMAN, G.A.; REZNITSKIY, L.A.; KOLESOV, V.P.;
GAL'CHENKO, G.L.; KHODEYEV, Yu.S.; KHACHKURUZOV, G.A.;
SOKOLOV, V.B.; GOROKHOV, L.N.; MONAYENKOVA, A.S.;
KOMAROVA, A.F.; VEYTS, I.V.; YURKOV, G.N.; MALENKOV, G.G.;
SMIRNOVA, N.L.; GLUSHKO, V.P., akademik, otv. red.;
MIKHAYLOV, V.V., red.; KARAPET'YANTS, M.Kh., red.

[Thermal constants of substances; reference book in ten
numbers] Termicheskie konstanty veshchestva; spravochnik
v desiati vypuskakh. Moskva, No.1. 1965. 144 p.
(MIRA 18:7)

1. Moscow. Vsesoyuznyy institut nauchnoy i tekhnicheskoy
informatsii.

U.S. 100-7-65 EPR(c)/EPR/EWT(j)/EWT(1)/EWT(m)/EWT(n) Pc-L/Pr-L/Ps-L/Pl-L RPL
 8/0294/65/003/001/0033/0046
 AP5006461

AUTHOR: Gurvich, L. V.; Rtishcheva, K. P.

TITLE: Analytic representation of tabulated values of thermodynamic properties of gases

SOURCE: Teplofizika vysokikh temperatur, v. 3, no. 1, 1965, 33-46

TOPIC TAGS: thermodynamic property, ideal gas, entropy, enthalpy, specific heat, thermodynamic potential, analytic approximation

ABSTRACT: In view of certain difficulties involved in the use of tabulated thermodynamic characteristics of various substances, especially when it comes to interpolation, the authors consider the possibility of representing thermodynamic properties of substances in an ideal-gas state by equations of the type $A_1 + A_2 T + A_3 T^2 + A_4 T^3 + A_5 T^4 + A_6 T^5 + A_7 T^6 + A_8 T^7 + A_9 T^8 + A_{10} T^9 + A_{11} T^{10}$ which would be convenient for use in electronic computers with limited memory capacity. It is shown that for most gases satisfactory accuracy can be obtained in the temperature ranges 293 - 6000K and 1000 - 20000K using $+2 \leq n \leq +3$. The coefficients of the corresponding equations are calculated for the 355 gases listed in

Card 1/2

L 45637-65

ACCESSION NR: AF5006467

a book by one of the authors (Gurvich et al, Termodinamicheskiye svoystva individual'nykh veshchestv [Thermodynamic Properties of Individual Substances], Izd-vo AN SSSR, 1962). The results are listed in tables. "In conclusion the authors thank B. A. Vorb'yev and V. S. Yungman for valuable discussion of the questions considered in this paper, and also I. G. Baybuz and V. S. Shmelova for help with the computer calculations." Orig. art. has: 15 formulas, 2 tables, and 1 figure.

ASSOCIATION: Nauchno-issledovatel'skiy institut vysokikh temperatur (Scientific Research Institute of High Temperatures)

SUBMITTED: 30May64

ENCL: 00

SUB CODE: TD,ME

NR REF SOV: 002

OTHER: 007

Card 2/2 mb

RYABOVA, V.G.; GURVICH, I.V.

Metal - hydroxyl bonding energy in CaOH, SrOH, and BaOH molecules.
Teplofiz. vys. temp. 3 no.2:318-321 Mr-Apr '65. (MIRA 18:7)

1. Nauchno-issledovatel'skiy institut vysokikh temperatur, Moskva.

L 21171-65 ENT(m)/ENP(t)/ENP(b) IJP(o) JD/JW
 ACCESSION NR: AP5003033

S/0051/65/018/001/0132/0134

AUTHOR: Gurvich, L. V.; Novikov, M. M.; Ryabova, V. G.

TITLE: Investigation of spectra and determination of dissociation energies of oxygen compounds of gallium and indium

SOURCE: Optika i spektroskopiya, vo. 18, no. 1, 1965, 132-134

TOPIC TAGS: arc spectrum, dissociation energy, gallium compound, indium compound, oxide, hydroxide

ABSTRACT: In view of the contradictory published data concerning the dissociation energy of the GaO and InO molecules, and also concerning the oxygen compounds produced by gallium and indium in flames, the authors have undertaken new investigations of the electronic spectrum of GaO and the equilibrium reaction of Ga and In with the combustion products of flames of the type $aH_2 + bO_2 + cN_2 + dH_2O$ and $aCO + bO_2 + dH_2O$. The known system of bands of GaO was investigated in the 3350--4150 Å band with a grating spectrograph, and the constants of the molecule GaO were determined. An attempt to obtain the absorption spectrum of GaO in the range 3600--7000 Å with the arc and gas discharge exposed to a strong pulsed

Card 1/2

L 21171-65

ACCESSION NR: AP5003033

source has shown absorption due to GaO only in the 4000 Å region. Although there is no final proof that the lower state of the investigated system is the ground state of GaO, this assumption is quite likely. The equilibrium of the reactions of Ga and In with the flame combustion products was investigated by determining the partial pressures of the metals from the relative intensity of the atomic lines in the flame spectra. The corresponding equilibrium constants were calculated from the measured partial pressures and from those calculated theoretically for equilibrium conditions. It was impossible to determine the dissociation energies of GaOH and InOH in the flames of carbon monoxide, because of the high temperature and the low concentration of the hydroxyl. In hydrogen and oxygen flames, the dissociation energies of GaOH and InOH were 101 ± 5 and 90 ± 5 kcal/mole, respectively. It is concluded that the main compounds of Ga and In in the 12 types of flames employed are the hydroxides, produced in the reaction $Me + H_2O = MeOH + H$ (Me = Ga or In). Other effects observed in the flames are briefly discussed.

ASSOCIATION: None

SUBMITTED: 04Nov63

ENCL: 00

SUB CODE: 00, *LC*

NR REF SOV: 003

OTHER: 004

Card 2/2

L 22894-65 EWT(m)/EWP(t)/EWP(b) IJP(c) JD

ACCESSION NR: AP5003037

S/0051/69/018/001/0143/0145

AUTHOR: Gurvich, L. V.; Ryabova, V. G.

TITLE: Investigation of the dissociation energy of BaO and BaOH

SOURCE: Optika i spektroskopiya, v. 18, no. 1, 1965, 143-145

TOPIC TAGS: barium compound, dissociation energy, binding energy, flame spectroscopy

ABSTRACT: In a book by one of the authors (with G. A. Khachkurazov, V. A. Medvedev, and I. V. Veyts "Termodinamicheskiye svoystva individual'nykh veshchestv" [Thermodynamic Properties of Individual Substances], AN SSSR, M., 1962) it is stated that the value of the dissociation energy D_0 obtained as a result of the investigation of equilibrium reactions of Ba in flames and by determining the heat of sublimation of BaO by various methods are in agreement, and yield a value 137 ± 2 kcal/mole. However, since these two methods of measurement can result in a considerable dispersion of the results, the authors have redetermined the dissociation energy by measuring the partial pressure of atomic barium in a flame of carbon monoxide with oxygen, in which no BaOH molecules are formed to distort

Card 1/2

L 22894-65

ACCESSION NR: AP5003037

the results. The measurements were made in the flame with composition $200 + O_2 + 0.03 H_2O$ ($T = 2,965K$). The partial pressure was determined from the absolute intensity of the $\lambda = 5535 \text{ \AA}$ line. At the same time, the partial pressure of the Ba^+ ions was determined from the absolute intensities of the 4,554 and 4,934 \AA lines. Check measurements were also made in three hydrogen-air flames with different compositions, and the results were much higher dissociation values, the difference being due to the fact that the flame contained appreciable amounts of $BaOH$. The effect of the presence of $BaOH$ on other measurements is discussed briefly, and the values of 134 ± 8 and $114 \pm 5 \text{ kcal/mole}$ respectively are recommended for the dissociation energies of the molecules BaO and for the binding energy of $BaOH$, respectively. Orig. art. has: 2 formulas and 1 table.

ASSOCIATION: None

SUBMITTED: 03Feb64

ENCL: 00

SUB CODE: 0P

NR REF SOV: 010

OTHER: 002

Card 2/2

L 4386-66 EWT(1)/T IIP(c)

ACC NR: AP5017909

UR/0051/65/019/001/0143/0145
535.33

AUTHOR: Novikov, M. M.; Gurvich, L. V.

TITLE: A new study of the emission spectrum of the SrCl molecule

SOURCE: Optika i spektroskopiya, v. 19, no. 1, 1965, 143-145

TOPIC TAGS: strontium compound, chloride, emission spectrum, band spectrum, optic transition, dissociation constant

ABSTRACT: One of the aims of the study was to obtain in emission the same systems of bands that were previously observed in absorption only, and thereby determine more accurately the constants of the molecule under investigation. The radiation source was an uncondensed discharge in a tube (H. Schuller, Spectrochim. Acta v. 4, 85, 1950), with an electrically heated capillary (140 mm long and 5 mm in diameter). The discharge voltage and current were 3--4 kv and 1.5 - 2 a. The spectra were photographed with a high-transmission spectrograph with STE-1 diffraction grating. The emission spectrum was found to contain all six previously known band systems, as well as additional bands, including some with vibrational quantum numbers larger than those reported earlier. The dissociation energies, frequencies, and vibrational constants of the transitions are calculated. Orig. art. has: 5 formulas and 1 table.

ASSOCIATION: None

Card 1/2